

Influence of thermophoresis and thermal radiation on mass transfer and heat of Three Dimensional maxwell fluid in presence of Magnetic field

Nidhal .M. AbdAlameer

nidhaalm_math@csw.uobaghdad.edu.iq

Department of mathematics , college of science for women , university of Baghdad , Baghdad , Iraq.

ABSTRACT

The study is about Maxwell , three dimensions of non – Newtonian fluid. Method of th Homotopy applied to analysis mass transfer and heat with thermophoresis effects. (Sc), Impact of therrmophoretic (τ), magnetic (M), Biot (γ), radiation (Rd), Schmidt Prandtle (Pr) parameters and ratio parameter(β) on concentration, temperature are offered in the paper.

Keywords:

Maxwell fluid, magnetic field, mass transfer , three dimensions fluid

1. Introduction

mass transfer, heat are processes submitted by mathematical equations in diffusion and convection ,This subject in the boundary flow is significant, for example, growing of crystal, , extrusion of polymer, glass-fiber, and many others . The phenomenon of movmment of fluid particles from warm region in the direction the cold region is known a thermophoresis, this motion occurs because molecules with higher kinetic energy from the warmer region , hit with molecules in cooler area with low energy.

A thermophoretic velocity mean the velocity gained the particles , while the force to which the particles are exposed are denominated the thermophoretic force . [1] .lately, researchers has been centering on model of Maxwell fluid.

Ganeshkumar.k,Rudraswamy.N.G,Gireesha.B.Ja nd Krishnamurth.M [2], made an impact for

viscous dissipation and thermal radiation of nano fluid with joule heating , method of Runge - kutta and shooting used to find the solution. Koneril and others [3], discussed in the attendance of dust, particles the impact of radiation , use Maxwell fluid with magnetic field in this paper used a numerical method to obtain the solution and it show that the effect of the parameters which appear on the flow . Must and others [4] submitted Maxwell fluid for sakiads flow with magnetic field . Vijayaa. N ,Krishna Jyothb. P , Anupamac. A , Leelavathid. R, Ambicae. K[6] presented model of Cattaneo- christov heat flux to show influence thermophorsis and buoyancy on chemical reaction of Maxwell fluid. This article discusses the impact of thermophoresis ,thermal radiation of Maxwell fluid with magnetic field in three dimensions space. The technique that was used to solve the problem is homotopy [5,8].

2. Formulation

Assume that the coordinates are Cartesian, three dimensional maxwell fluid, the field is magnetic, the surface is inclined stretching, T_∞

is temperature interface and T_f is surface temperature. The equations for present flow are

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = 0 \tag{1}$$

$$u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} + w \frac{\partial u}{\partial z} = \nu \frac{\partial^2 u}{\partial y^2} - \lambda \left(u^2 \frac{\partial^2 u}{\partial x^2} + v^2 \frac{\partial^2 u}{\partial y^2} + w^2 \frac{\partial^2 u}{\partial z^2} + 2(uv \frac{\partial^2 u}{\partial x \partial y} + vw \frac{\partial^2 u}{\partial y \partial z} + uw \frac{\partial^2 u}{\partial x \partial z}) \right)$$

$$\mu - \frac{\sigma \beta_0^2}{\rho} \tag{2}$$

$$u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} + w \frac{\partial v}{\partial z} = \nu \frac{\partial^2 v}{\partial y^2} - \lambda \left(\frac{u^2 \partial^2 v}{\partial x^2} + \frac{v^2 \partial^2 v}{\partial y^2} + \frac{w^2 \partial^2 v}{\partial z^2} + 2(uv \frac{\partial^2 v}{\partial x \partial y} + vw \frac{\partial^2 v}{\partial y \partial z} + wu \frac{\partial^2 v}{\partial x \partial z}) \right) \tag{3}$$

$$\rho c_p \left(u \frac{\partial T}{\partial x} + v \frac{\partial T}{\partial y} + w \frac{\partial T}{\partial z} \right) = \frac{\partial}{\partial z} \left(\left(\frac{16\sigma^*}{3k^*} T_\infty^3 + k \right) \frac{\partial T}{\partial z} \right) \tag{4}$$

$$u \frac{\partial c}{\partial x} + v \frac{\partial c}{\partial y} + w \frac{\partial c}{\partial z} = D_e \frac{\partial^2 c}{\partial z^2} - \frac{\partial}{\partial z} (V_T (c - c_\infty)) \tag{5}$$

Considering that

v, u with w represent the velocities

λ relaxation time

σ^* electrical conductivity

g the gravitational acceleration

T mean the temperature

β_T thermal coefficient

β_c concentration expansion coefficient

$\nu = \left(\frac{\mu}{\rho} \right)$ kinematic viscosity, μ, ρ are represent the dynamic viscosity and density of fluid respectively

σ thermal diffusivity

C concentration

K_T ratio of thermal diffusion

$V_T = -k_2 \frac{\nu}{T_r} \frac{\partial T}{\partial z}$ thermophoretic velocity,

k_2 thermophoretic coefficient

T_r reference temperature

D_e mass diffusivity

$\tau = -k_2 \frac{T_f - T_\infty}{T_r}$ Thermophoretic parameter

β_0 Strength of Magnetic field

k thermal conductivity

T_f ambient temperature

C_∞ ambient concentration

Put the boundary conditions

$$u = u_e = \alpha x$$

$$v = \Gamma_y, w = 0, -k^*/(T_f - T_\infty) \frac{\partial T}{\partial z} = h, \quad c = c_w \quad \text{at } z = 0$$

u converge to zero, v converge to zero, T converge to T_∞ , C converge to C_∞ at $z \rightarrow \infty$

Note that α and Γ owns dimension reverse time.

Let

$$\left. \begin{aligned} u &= \alpha_x f'(\eta) \\ v &= \alpha_y g'(\eta) \\ w &= -\sqrt{\alpha v} (f(\eta) + g(\eta)) \end{aligned} \right\} \quad (6)$$

$$\theta(\eta) = (T - T_\infty) / (T_f - T_\infty)$$

$$\eta = z \sqrt{\alpha / v}$$

$$\Phi(\eta) = (c - c_\infty) / (c_w - c_\infty)$$

Substitute (6) in equations(2- 5)

$$f''' + (f + g) f'' - M f' - \beta_1 [(f + g)^2 f''' - 2f'(f + g) f''] - (f')^2 = 0 \quad (7)$$

$$g''' + (f + g) g'' + \beta_1 [2(f + g) g'g'' - (f + g)^2 g'''] - (g')^2 = 0, \quad (8)$$

$$(1 + 4R/3) \theta'' + pr \theta' (f + g) = 0 \quad (9)$$

$$\Phi'' + Sc \Phi' (f + g) - Sc\tau (\Phi' \theta' - \Phi \theta'') = 0 \quad (10)$$

and

at $\eta = 0$

$$f = 0$$

$$f' = 1$$

$$g = 0$$

$$g' = \beta$$

$$\theta' = -\gamma(1 - \theta(0)), \quad \Phi = 1$$

at η converge to ∞

f' converge to 0

g' converge to 0

θ converge to 0

ϕ converge to 0

Note $\beta_1 = \lambda_1 a$, $\beta = b/a$, $v\sigma = pr$, $\gamma = h/k^* \sqrt{v/a}$, (biot number), σ^* represent Stefan -Boltzmann constant, β_T is the coefficients of thermal expansion, c_p mean specific heat, k^* the coefficient of absorption, $k = k_1 a$, $sc = v/D$, $\lambda = GrxRe2x$ (local buoyancy), $Grx = g$

$\beta_T (T_f - T_\infty) \frac{x^3}{\nu^2}$ (local grashof number) , $\frac{\beta C (C_w - C_\infty)}{\beta_T (T_f - T_\infty)} = N$ (constant dimensionless concentration buoyancy parameter) , $M = \frac{\sigma \beta_0^2}{a \rho}$ (parameter of magnetic field) .

3. Solution method

To resolve the issue ,we'll use homotopy method , which demands the approximations

$$\begin{aligned} f_0(\eta) &= 1 - \exp(-\eta) \\ g_0(\eta) &= -(\exp(-\eta) - 1) \beta \\ \theta_0(\eta) &= (\exp(-\eta) + 1) \gamma \\ \phi_0(\eta) &= \exp(-\eta) \end{aligned}$$

and the linear transformations

$$L_1 = f''''(\eta) - f'(\eta) , L_2 = g''''(\eta) - g'(\eta)$$

$$L_3 = \theta''(\eta) - \theta(\eta) , L_4 = \Phi''(\eta) - \Phi(\eta),$$

$$L_1 (A_1 + A_2 e^\eta + A_3 e^{-\eta}) = 0,$$

$$L_2 (A_4 + A_5 e^\eta + A_6 e^{-\eta}) = 0$$

$$L_3 (A_7 e^\eta + A_8 e^{-\eta}) = 0,$$

(11)

$$L_4 (A_9 e^\eta + A_{10} e^{-\eta}) = 0,$$

$A_1, A_2, A_3, \dots, A_{10}$ are constants.

The issue is from order zero :

$$(1-\mathfrak{z}) L_1 [\tilde{f}(\eta; \mathfrak{z}) - f_0(\eta)] = \mathfrak{z} \hbar_1 N_1 [\tilde{f}(\eta; \mathfrak{z}), \tilde{g}(\eta; \mathfrak{z})],$$

$$(1-\mathfrak{z}) L_2 [\tilde{g}(\eta; \mathfrak{z}) - g_0(\eta)] = \mathfrak{z} \hbar_2 N_2 [\tilde{f}(\eta; \mathfrak{z}), \tilde{g}(\eta; \mathfrak{z})],$$

$$(1-\mathfrak{z}) L_3 [\tilde{\theta}(\eta; \mathfrak{z}) - \theta_0(\eta)] = \mathfrak{z} \hbar_1 N_1 [\tilde{f}(\eta; \mathfrak{z}), \tilde{g}(\eta; \mathfrak{z}), \tilde{\theta}(\eta; \mathfrak{z}), \tilde{\Phi}(\eta; \mathfrak{z})],$$

$$(1-\mathfrak{z}) L_4 [\tilde{\Phi}(\eta; \mathfrak{z}) - \Phi_0(\eta)] = \mathfrak{z} \hbar_2 N_2 [\tilde{f}(\eta; \mathfrak{z}), \tilde{g}(\eta; \mathfrak{z}), \tilde{\theta}(\eta; \mathfrak{z}), \tilde{\Phi}(\eta; \mathfrak{z})],$$

$$\tilde{f}(0, \mathfrak{z}) = 0, \tilde{f}'(\infty, \mathfrak{z}) = 0, \tilde{f}'(0, \mathfrak{z}) = 1,$$

$$\tilde{g}(0, \mathfrak{z}) = 0, \tilde{g}'(\infty, \mathfrak{z}) = 0, \tilde{g}'(0, \mathfrak{z}) = \beta, \tilde{g}(\infty, \mathfrak{z}) = 0$$

$$\tilde{\theta}'(0, \mathfrak{z}) = -\gamma [1 - \theta(0, \mathfrak{z})], \tilde{\theta}(\infty, \mathfrak{z}) = 0, \tilde{\Phi}(0, \mathfrak{z}) = 1, \tilde{\Phi}(\infty, \mathfrak{z}) = 0$$

$$N_1[\tilde{f}(\eta; \zeta), \tilde{g}(\eta; \zeta)] = \frac{\partial^3 \tilde{f}(\eta; \zeta)}{\partial \eta^3} + (\tilde{f}(\eta; \zeta) + \tilde{g}(\eta; \zeta)) \frac{\partial^2 \tilde{f}(\eta; \zeta)}{\partial \eta^2} - M \frac{\partial \tilde{f}(\eta; \zeta)}{\partial \eta} + \beta^* \left[\frac{\partial^2 \tilde{f}(\eta; \zeta)}{\partial \eta} \right. \\ \left. (\tilde{f}(\eta; \zeta) + \tilde{g}(\eta; \zeta)) \frac{\partial^2 \tilde{f}(\eta; \zeta)}{\partial \eta^2} - (\tilde{f}(\eta; \zeta) + \tilde{g}(\eta; \zeta))^2 \frac{\partial^3 \tilde{f}(\eta; \zeta)}{\partial \eta^3} \right] - \left(\frac{\partial \tilde{f}(\eta; \zeta)}{\partial \eta} \right)^2 \quad (12)$$

$$N_2[\tilde{f}(\eta; \zeta), \tilde{g}(\eta; \zeta)] = \frac{\partial^3 \tilde{g}(\eta; \zeta)}{\partial \eta^3} + (\tilde{f}(\eta; \zeta) + \tilde{g}(\eta; \zeta)) \frac{\partial^2 \tilde{g}(\eta; \zeta)}{\partial \eta^2} + \beta^* [2(\tilde{f}(\eta; \zeta) + \tilde{g}(\eta; \zeta)) \frac{\partial \tilde{g}(\eta; \zeta)}{\partial \eta} \\ \frac{\partial^2 \tilde{g}(\eta; \zeta)}{\partial \eta^2} - (\tilde{f}(\eta; \zeta) + \tilde{g}(\eta; \zeta))^2 \frac{\partial^3 \tilde{g}(\eta; \zeta)}{\partial \eta^3}] - \left(\frac{\partial \tilde{g}(\eta; \zeta)}{\partial \eta} \right)^2 \quad (13)$$

$$N_3[\tilde{\theta}(\eta, \zeta), \tilde{\Phi}(\eta, \zeta), [\tilde{f}(\eta, \zeta), \tilde{g}(\eta, \zeta)]] = (1 + 4/3 R) \frac{\partial^2 \tilde{\theta}(\eta, \zeta)}{\partial \eta^2} + pr(\tilde{f}(\eta, \zeta) + \tilde{g}(\eta, \zeta)) \\ \left. \right) \frac{\partial \tilde{\theta}(\eta, \zeta)}{\partial \eta} \quad (14)$$

$$N_4[\tilde{\Phi}(\eta, \zeta), \tilde{\theta}(\eta, \zeta), \tilde{f}(\eta, \zeta), \tilde{g}(\eta, \zeta)] = \frac{\partial^2 \tilde{\Phi}}{\partial \eta^2} + sc(\tilde{f}(\eta, \zeta) + \tilde{g}(\eta, \zeta)) \frac{\partial \tilde{\Phi}}{\partial \eta} - sc\tau \left(\frac{\partial \tilde{\Phi}}{\partial \eta} \frac{\partial \tilde{\theta}(\eta, \zeta)}{\partial \eta} \right. \\ \left. - \tilde{\Phi}(\eta, \zeta) \frac{\partial^2 \tilde{\theta}(\eta, \zeta)}{\partial \eta^2} \right), \quad (15)$$

ζ and \hbar_1, \dots, \hbar_n ($n=1,2,3,4$) mean embedding , auxiliary parameters respectively

N_1, N_2, N_3, N_4 nonlinear transformations.

at $\zeta = 0, \zeta = 1$

$$\tilde{f}(\eta, 0) = f_0(\eta), \tilde{f}(\eta, 1) = f(\eta), \tilde{g}(\eta, 0) = g_0(\eta), \tilde{g}(\eta, 1) = g(\eta), \\ \tilde{\theta}(\eta, 0) = \theta_0(\eta), \tilde{\Phi}(\eta, 0) = \phi_0(\eta), \tilde{\theta}(\eta, 1) = \theta(\eta), \tilde{\Phi}(\eta, 1) = \phi_0(\eta). \quad (16)$$

By Taylor's expansion

$$f(\eta, \zeta) = f_0(\eta) + \sum_{m=1}^{\infty} f_m(\eta) \zeta^m, f_m(\eta) = 1/(m!) \left[\frac{\partial^m f(\eta; \zeta)}{\partial \eta^m} \right]_{\zeta=0} \\ g(\eta, \zeta) = g_0(\eta) + \sum_{m=1}^{\infty} g_m(\eta) \zeta^m, g_m(\eta) = 1/(m!) \left[\frac{\partial^m g(\eta; \zeta)}{\partial \eta^m} \right]_{\zeta=0} \quad (17)$$

$$\theta(\eta, \zeta) = \theta_0(\eta) + \sum_{m=1}^{\infty} \theta_m(\eta) \zeta^m, \theta_m(\eta) = 1/(m!) \left[\frac{\partial^m \theta(\eta; \zeta)}{\partial \eta^m} \right]_{\zeta=0}$$

$$\phi(\eta, \zeta) = \phi_0(\eta) + \sum_{m=1}^{\infty} \phi_m(\eta) \zeta^m, \phi_m(\eta) = 1/(m!) \left[\frac{\partial^m \phi(\eta; \zeta)}{\partial \eta^m} \right]_{\zeta=0}$$

by choosing the series (17) convergent when $\zeta=1$ so

$$f(\eta) = f_0(\eta) + \sum_{m=1}^{\infty} f_m(\eta),$$

$$g(\eta) = g_0(\eta) + \sum_{m=1}^{\infty} g_m(\eta),$$

$$\theta(\eta) = \theta_0(\eta) + \sum_{m=1}^{\infty} \theta_m(\eta),$$

$$\phi(\eta) = \phi_0(\eta) + \sum_{m=1}^{\infty} \phi_m(\eta).$$

The issue that's generated from deformation of order m :

$$L_1 [f_m (\eta) - x_m f_{m-1} (\eta)] = \hbar_1 R_{1m} (\eta) ,$$

$$L_2 [g_m (\eta) - x_m g_{m-1} (\eta)] = \hbar_2 R_{2m} (\eta) ,$$

$$L_3 [\theta_m (\eta) - x_m \theta_{m-1}(\eta)] = \hbar_3 R_{3m} (\eta) ,$$

$$L_4 [\Phi_m (\eta) - x_m \Phi_{m-1}(\eta)] = \hbar_4 R_{4m} (\eta) ,$$

$$f_m (0) = 0 = g_m (0) \quad , \quad f'_m (0) = g'_m (0) = 0 = f'_m (\infty) = g'_m (\infty)$$

$$\theta'_m (0) - \gamma \theta_m (0) = 0$$

$$\theta_m (\infty) = 0 = \Phi_m (\infty) = \Phi_m (0)$$

$$R_1^m = D[f_{-1+m} , \eta , \eta , \eta] - (M * D[f_{-1+m} , \eta]) - \sum_{k=0}^{m-1} ([f_{m-1-k} , \eta] * D[f_k, \eta]) + \sum_{k=0}^{m-1} ((f_{m-1-k} D[f_k, \eta, \eta]) + \beta \sum_{k=0}^{m-1} \sum_{i=0}^k ((f_{m-1-k} + g_{m-1-k}) * 2D[f_{k-i}, \eta] * D[f_i, \eta, \eta]) - ((((f_{m-1-k}) (f_{k-i})) + ((g_{m-1-k}) (g_{k-i}))) + (((f_{m-1-k}) (g_{k-i})))) * 2D[f_i, \eta, \eta] .$$

(18)

$$R_2^m = D[g_{m-1}, \eta, \eta, \eta] - \sum_{k=0}^{m-1} (D[g_{m-k-1}, \eta] D[g_k, \eta]) + \sum_{k=0}^{m-1} ((f_{m-k-1} D[g_k, \eta, \eta]) + ((g_{m-1-k} D[g_k, \eta, \eta])) + \beta$$

$$\sum_{k=0}^{m-1} \sum_{i=0}^k (2 D[g_{k-i}, \eta] (f_{m-1-k} + g_{m-k-1}) D[g_i, \eta, \eta] - (((f_{m-k-1} f_{k-i}) + (g_{m-k-1} g_{k-i}) + (2 * (f_{m-k-1} g_{k-i}))) D[g_i, \eta, \eta , \eta] .$$

(19)

$$R_3^m = (1 + (4/3)Rd) * D[\theta_{m-1}, \eta, \eta] + (Pr * \sum_{k=0}^{m-1} ((D[\theta_{m-1-k}, \eta] * f_k) + (D[\theta_{m-1-k}, \eta] * g_k)))$$

(20)

$$R_4^m = D[\Phi_{m-1}, \eta, \eta] + (Sc * \sum_{k=0}^{m-1} ((D[\Phi_{m-1-k}, \eta] * f_k + (D[\Phi_{m-1-k}, \eta] * g_k))) - (Sc * \tau * \sum_{k=0}^{m-1} ((D[\Phi_{m-1-k}, \eta]) *$$

$$D[\theta_k, \eta] - (\Phi_{m-1-k} * D[\theta_k, \eta, \eta])))$$

..(21)

It easy to solve the equations(16-19) using mathematic software ,so we get

$$f_1 = \frac{1}{24} e^{-3\eta} (h_1 \beta (-1 + \beta^2) - 4e^\eta h_1 \beta (1 + 2\beta + 2\beta^2) - 6e^{2\eta} (h_1 (M + \beta(2 + 2 + \beta^2)) (3 + 2\eta) + 4(\frac{1}{24} (24 - 6h_1 M - 23h_1 \beta - 28h_1 \beta^2 - 19h_1 \beta^3))))) .$$

(22)

$$f_2 = \frac{1}{2880} e^{-5\eta} (h^2 \beta^2 (3 + 32\beta + 22\beta^2 - 32\beta^3 - 25\beta^4) + 8e^\eta h^2 \beta (-1 - 14\beta - 20\beta^2 + 13\beta^3 + 53\beta^4 + 29\beta^5) + 5e^{2\eta} h \beta (96(-1 + \beta^2) + h (-26 + 9\beta^5 (-15 + 4\eta) + 4\beta^3 (-80 + 9\eta) + 8\beta^4 (-43 + 9\eta) - 2\beta^2 (91 + 36\eta) - \beta (73 + 72\eta) + 2M(-18(1 + \eta) + \beta^2 (-7 + 6\eta)))))) \quad ..(23)$$

$$f_3 = \frac{1}{2419200} e^{-7\eta} (5h^3 \beta^3 (-155 - 1408\beta - 3967\beta^2 - 2880\beta^3 + 2695\beta^4 + 4288\beta^5 + 1427\beta^6) - 4e^\eta h^3 \beta^2 (-240 - 4751\beta - 24522\beta^2 - 39788\beta^3 + 36\beta^4 + 67749\beta^5 + 67862\beta^6 + 20054\beta^7) - 7e^{2\eta} h^2 \beta (840\beta (-3 - 32\beta - 22\beta^2 + 32\beta^3 + 25\beta^4)) \quad ..(24)$$

$$f_4 = -\frac{1}{174182400} e^{-9\eta} h^4 \beta^4 (-17619 - 213424\beta - 917030\beta^2 - 1745360\beta^3 - 1162248\beta^4 + 859184\beta^5 + 1874270\beta^6 + 1099600\beta^7 + 222627\beta^8) + \frac{1}{304819200} e^{-8\eta} h^4 \beta^3 (-41910 - 1003527\beta - 7123122\beta^2 - 22075329\beta^3 - 30357442\beta^4 - 4581577\beta^5 + 40482514\beta^6 + 52695097\beta^7 + 27280216\beta^8 + 5205080\beta^9) \quad ..(25)$$

$$g_1 = \frac{1}{24} e^{-3\eta} (-h^2 \beta^2 (-1 + \beta^2) - 4e^\eta h^2 \beta (1 + 2\beta + 2\beta^2) - 6e^{2\eta} (h^2 \beta^2 (2 + 2\beta + \beta^2) (3 + 2\eta) + 4(\frac{1}{24} (24\beta - 8h^2 \beta - 25h^2 \beta^2 - 28h^3 \beta^3 - 9h^2 \beta^4)))) + \frac{1}{12} (12\beta - 2h^2 \beta + 9h^2 \beta^2 + 8h^2 \beta^3 + 5h^2 \beta^4) \quad ..(26)$$

$$g_2 = \frac{1}{2880} e^{-5\eta} (h^2 \beta^3 (-25 - 32\beta + 22\beta^2 + 32\beta^3 + 3\beta^4) - 8e^\eta h^2 \beta^2 (-10 - 38\beta - 47\beta^2 - 2\beta^3 + 29\beta^4 + 8\beta^5) - 5e^{2\eta} h^2 \beta (96\beta (-1 + \beta^2) + h^2 (16 + \beta^2 (265 - 72\eta) + \beta^3 (386 - 72\eta) + 12\beta^4 (23 + 3\eta) + 8\beta^5 (5 + 9\eta) + \beta^6 (3 + 36\eta) + \beta (94 - 50M - 24M\eta))) \quad ..(27)$$

$$g_3 = \frac{1}{2419200} e^{-7\eta} (-5h^2 \beta^3 (-1427\beta^4 - 4288\beta^5 - 2695\beta^6 + 2880\beta^7 + 3967\beta^8 + 1408\beta^9 + 155\beta^6) + 4e^\eta h^2 \beta^3 (-5361\beta^3 - 34054\beta^4 - 73844\beta^5 - 57204\beta^6 + 16603\beta^7 + 46010\beta^8 + 19338\beta^9 + 2112\beta^{10})) \quad ..(28)$$

$$\theta_1 = \frac{1}{12(1+\gamma)} e^{-2\eta} h \gamma (-3e^\eta + 4Pr + 3e^\eta Pr - 4e^\eta Rd + 4Pr\beta + 3e^\eta Pr\beta - 6e^\eta \eta + 6e^\eta Pr\eta - 8e^\eta Rd\eta + 6e^\eta Pr\beta\eta) + e^{-\eta} ((12\gamma - 3h\gamma - 5hPr\gamma - 4hRd\gamma - 5hPr\beta\gamma + 12\gamma^2)/(12(1 + \gamma))) \quad ..(29)$$

$$\theta_2 = \frac{1}{2160(1+\gamma)} e^{-4\eta} (-1080e^{3\eta} h \gamma - 135e^{3\eta} h^2 \gamma + 2160e^{2\eta} hPr\gamma + 1620e^{3\eta} hPr\gamma + 480e^{2\eta} h^2 Pr\gamma + 360e^{3\eta} h^2 Pr\gamma + 840e^{2\eta} h^2 MPr\gamma + 270e^{3\eta} h^2 MPr\gamma + 180e^\eta h^2 Pr^2 \gamma - 480e^{2\eta} h^2 Pr^2 \gamma - 225e^{3\eta} h^2 Pr^2 \gamma - 1440e^{3\eta} hRd\gamma - 360e^{3\eta} h^2 Rd\gamma + 640e^{2\eta} h^2 PrRd\gamma + 480e^{3\eta} h^2 PrRd\gamma - 240e^{3\eta} h^2 Rd^2 \gamma + 2160e^{2\eta} hPr\beta\gamma + 1620e^{3\eta} hPr\beta\gamma + 6h^2 Pr\beta\gamma + 90e^\eta h^2 Pr\beta\gamma + 1590e^{2\eta} h^2 Pr\beta\gamma + 675e^{3\eta} h^2 Pr\beta\gamma + 360e^\eta h^2 Pr^2 \beta\gamma - 960e^{2\eta} h^2 Pr^2 \beta\gamma - 450e^{3\eta} h^2 Pr^2 \beta\gamma + 640e^{2\eta} h^2 PrRd\beta\gamma + 480e^{3\eta} h^2 PrRd\beta\gamma \quad ..(30)$$

$$\theta_3 = \frac{1}{5443200(1+\gamma)} e^{-6\eta} (-4082400e^{5\eta} h \gamma - 1020600e^{5\eta} h^2 \gamma - 170100e^{5\eta} h^3 \gamma + 10886400e^{4\eta} hPr\gamma + 8164800e^{5\eta} hPr\gamma + 6955200e^{4\eta} h^2 Pr\gamma + 4309200e^{5\eta} h^2 Pr\gamma + 957600e^{4\eta} h^3 Pr\gamma + 1058400e^{5\eta} h^3 Pr\gamma + 8467200e^{4\eta} h^2 MPr\gamma + 2721600e^{5\eta} h^2 MPr\gamma + 3427200e^{4\eta} h^3 MPr\gamma +$$

$$1512000e^{5\eta}h^3MPr\gamma + 2242800e^{4\eta}h^3M^2Pr\gamma + 510300e^{5\eta}h^3M^2Pr\gamma + 2268000e^{3\eta}h^2Pr^2\gamma - 6048000e^{4\eta}h^2Pr^2\gamma - 2835000e^{5\eta}h^2Pr^2\gamma + 756000e^{3\eta}h^3Pr^2\gamma - 1915200e^{4\eta}h^3Pr^2\gamma - 1152900e^{5\eta}h^3Pr^2\gamma + 982800e^{3\eta}h^3MPr^2\gamma - 2520000e^{4\eta}h^3MPr^2\gamma - 945000e^{5\eta}h^3MPr^2\gamma + 90720e^{2\eta}h^3Pr^3\gamma - 415800e^{3\eta}h^3Pr^3\gamma + 504000e^{4\eta}h^3Pr^3\gamma + 264600e^{5\eta}h^3Pr^3\gamma - 5443200e^{5\eta}hRd\gamma .$$

..(31)

$$\phi_1 = \frac{1}{12}e^{-2\eta}h(-3e^\eta + 4Sc + 3e^\eta Sc + 4Sc\beta + 3e^\eta Sc\beta - 6e^\eta\eta + 6e^\eta Sc\eta + 6e^\eta Sc\beta\eta) + e^{-\eta}(\frac{1}{12}(12 + 3h - 7hSc - 7hSc\beta)).$$

..(32)

$$\phi_2 = \frac{1}{720(1+\gamma)}e^{-4\eta}(-360e^{3\eta}h - 135e^{3\eta}h^2 + 720e^{2\eta}hSc + 540e^{3\eta}hSc + 280e^{2\eta}h^2Sc + 240e^{3\eta}h^2Sc + 280e^{2\eta}h^2MSc + 90e^{3\eta}h^2MSc + 60e^\eta h^2Sc^2 - 200e^{2\eta}h^2Sc^2 - 105e^{3\eta}h^2Sc^2 + 720e^{2\eta}hSc\beta + 540e^{3\eta}hSc\beta + 2h^2Sc\beta + 30e^\eta h^2Sc\beta + 650e^{2\eta}h^2Sc\beta + 345e^{3\eta}h^2Sc\beta + 120e^\eta h^2Sc^2\beta - 400e^{2\eta}h^2Sc^2\beta - 210e^{3\eta}h^2Sc^2\beta - 2h^2Sc\beta^2 + 60e^\eta h^2Sc\beta^2).$$

..(33)

$$\phi_3 = \frac{1}{1814400(1+\gamma)^2}e^{-6\eta}(-1360800e^{5\eta}h - 1020600e^{5\eta}h^2 - 283500e^{5\eta}h^3 + 3628800e^{4\eta}hSc + 2721600e^{5\eta}hSc + 3528000e^{4\eta}h^2Sc + 2646000e^{5\eta}h^2Sc + 747600e^{4\eta}h^3Sc + 743400e^{5\eta}h^3Sc + 2822400e^{4\eta}h^2MSc + 907200e^{5\eta}h^2MSc + 1495200e^{4\eta}h^3MSc + 630000e^{5\eta}h^3MSc + 747600e^{4\eta}h^3M^2Sc + 170100e^{5\eta}h^3M^2Sc + 756000e^{3\eta}h^2Sc^2 - 2520000e^{4\eta}h^2Sc^2 - 1323000e^{5\eta}h^2Sc^2 + 327600e^{3\eta}h^3Sc^2 - 974400e^{4\eta}h^3Sc^2 - 585900e^{5\eta}h^3Sc^2 + 327600e^{3\eta}h^3MSc^2 - 974400e^{4\eta}h^3MSc^2 - 365400e^{5\eta}h^3MSc^2 + 30240e^{2\eta}h^3Sc^3 - 163800e^{3\eta}h^3Sc^3 + 243600e^{4\eta}h^3Sc^3 + 126000e^{5\eta}h^3Sc^3 + 3628800e^{4\eta}hSc\beta + 2721600e^{5\eta}hSc\beta + 30240e^{2\eta}h^2Sc\beta + 378000e^{3\eta}h^2Sc\beta$$

..(34)

Discussion and result

It's clear that all figure (1-11) are drawn for explain the influence the radiation parameter *Rd* and numbers Biot γ Prandtl pr , Schmidt *Sc*, stretching ratio η , Magnatic parameter *M*, thermophoretic parameter τ , on temperature concentration.

Effect γ, pr, β, Rd on temperature

Figs (1) illustrate the affected of parameter *Rd* on temperature. Instances $pr=0.7$ and the parameter $\beta=0.3, M=0.5$, we note that as *Rd* increase, the temperature increase. Figs (2) illustrate the temperature with impact parameter γ , its noted with Increase in γ the temperature increase.

Figs(3) show effect β when $\gamma=0.4, h=-1.4, Pr=0.7, Rd=0.3, M=0.5$ onto temperature

It's clear that the temperature is decrease with increase β . Observed in Fig (4) increase pr leads to decreased temperature.




Fig1: Influence the radiation parameter *Rd* on temperature when $\gamma=0.4, h=-1.4, pr=0.7, \beta=0.3, Rd=0, 0.3, 0.6, M=0.5$




Fig2: Influence the Biot parameter on temperature , $\gamma = 0.5, 0.7, 1, h = -1.4, pr = 0.7, \beta = 0.3, Rd = 0.3, M = 0.5$




Fig3: Influence the stretching ratio on temperature , $\gamma = 0.4, h = -1.4, Pr = 0.7, \beta = 0, 0.4, 0.6, Rd = 0.3, M = 0.5$




Fig4: Influence Péclet parameter on temperature , $\gamma = 0.4, h = -1.4, pr = 0.1, 0.5, 0.7, Rd = 0.3, \beta = 0.3, M = 0.5$

effect $\gamma, Rd, \tau, \beta, pr, M, Sc$ onto Concentration

All figures(5-11) indicated the concentration affected by parameters $\gamma, M, \beta, Rd, \tau, Pr, Sc$ when $\gamma = [0.5, 0.7, 1], \beta = [0.3, 0.7, 1],$

$pr = [1, 2, 3], Rd = [0.5, 0.7, 1], M = [0.5, 1, 2], \tau = [0.2, 0.5, 1], sc = [0, 0.5, 0.7]$. In Fig(5) its observed that at increase pr the concentration are

increase .Fig (6) its observed that with increase of Rd from 0.5 to 1 the concentration increase , when $\eta < 2$,For $\eta > 2$ the

concentration start to decrease. In Fig (7) the concentration are decrease with the variant sc from 0 to 0.7 . In fig 8 its noted that

when m increases ,the concentration decreases this similar in fig(7) .fig 9 explain effect β on the concentration when $\beta = 0.3, 0.7, 1$, its clear from the increase in β make the concentration increase. Fig10 is drawn to explain the effect τ on the concentration when $\tau = 0.2, 0.5, 1$, its noted that at τ increase the concentration decrease .In fig 11 effect γ on the concentration is similar to fig 10 .




Fig5: Influence Prandtl parameter on concentration when $\gamma=1, h=-1.4, Pr=1,2,3, \beta=0.5, Rd=1, M=1, \tau=1, Sc=1$




Fig6: Influence radiation parameter on concentration when $Rd=0.5,0.7, \gamma=0.7, h=-1.4, Pr=1, \beta=0.7, M=0.5, \tau=1, Sc=1$




Fig7: Influence Schmidt parameter on concentration when $\gamma=0.4, h=-1.4, Rd=0.3, \beta=0.3, pr=0.7, M=0.5, \tau=0.2, Sc=0.2,0.5,0.7$




Fig8: Influence magnetic parameter on concentration when $\gamma=1, h=-1.4, Pr=1, \beta=0.7, Rd=1, M=0.5,1.2, \tau=0.2, Sc=1$




Fig9: Influence stretching ratio parameter on concentration when $\gamma=1, h=-1.4, Pr=1, \beta=0.3,0.7,1, Rd=1, M=1.3, \tau=0.1, Sc=1$




Fig10: Influence thermophoretic parameter on concentration when $h=1.4, \gamma=1, Pr=0.5, \beta=1, M=1.3, \tau=0.2, 0.5, 1, Sc=1$




Fig11: Influence the Biot parameter on concentration when $\gamma=0.5, 0.7, 1, h=1.4, \mu=0.7, \beta=0.3, Rd=0.3, M=0.5, \tau=2, Sc=1$

References

1. Bilal .M , Sagheer. M, and Hussain . S . (2017) " On MHD 3D upper convected Maxwell fluid flow with thermophoretic effect using nonlinear radiative heat flux" , Publication: Canadian Journal of Physics <https://doi.org/10.1139/cjp-2017-0250> .
2. Ganeshkumar . k , Rudraswamy . N . G , Gireesha .B. J and Krishnamurth .M.R (2017) ." Influence of nonlinear thermal radiation and viscous dissipation on three-dimensional flow of Jeffrey nano fluid over a stretching sheet in the presence of Joule heating".journal nonlinear Engineering <https://doi.org/10.1515/nleng-2017-0014>.
3. koneril. K. s , Bijjanal J. G , Basavaja.M , Rama.S ."Influence of nonlinear thermal radiation and magnetic field on upper-convected Maxwell fluid flow due to aconvectively heat and stretching sheet in the presense of dust particles" .International Scientific publication and consulting services ,volume 2016, ISSUE117 .
4. Mustafa .M , Khan .J . A , Hayat .T , and Alsaedi. A. (2014)"Sakiadis flow of Maxwell fluid considering magnetic field and convective boundary conditions" , DOI : 10.1371/journal.pone.0103946 .
5. Nidhal.M.Abdal Ameer. (2014)." Influence of Thermal Radiation and MHD on the Boundary Layer Flow Due to an Exponentially Stretching Sheet", Iraqi Journal of Science, Vol 55, No.2A, pp:530-536
6. Vijayaa. N ,Krishna Jyothb. P., Anupamac. A , Leelavathid. R , Ambicae K . (2021) "Thermophoresis and Buoyancy Effects on Chemically Reactive Upper Convected Maxwell Fluid Flow Induced By An Exponentially Stretching Sheet: Application Of Cattaneo- Christov Heat Flux". Frontiers in Heat and Mass Transfer (FHMT), 17, 23 DOI: 10.5098/hmt.17.23
7. Raju. C.S.K., Sandeep. N, Sulochana.C, Sugnamma. V , JayachandraBabu. M. (2015) "Radiation, Inclined Vmagnetic field and cross-diffusion effects on flow over a stretching surface" .Journal of the Nigerian Mathematical Society 34(2015) 169-180.

-
8. Shijun. L .Beyond Perturbation . (2004)"Introduction to the Homotopy Analysis method ",Chapman and Hall/CRC press LLC. over an linearly stretching surface with convective condition Polson 2015:10(12) e0145332.
 9. Tasawar.H,Arslan.A,Taseer.M,Bashir.A. 2015"Influence of magnetic field in three dimensional flow of couple stress nanofluid