

Synthesis, characterization and antibacterial activities of new substituted (1,3)oxazepine 1,5-diones

Omar J. Mahdi Al-Asafi^{1,a)},

1) General Directorate of Education in Anbar, Ministry of Education, Iraq
a) Corresponding author Tel.: +964 -7812966986 Email: omerKha22@gmail.com, omar.j.mahdi.chem2021@st.tu.edu.iq

Mohammed G. Mukhlif^{1,b)}

1) General Directorate of Education in Anbar, Ministry of Education, Iraq
b) Corresponding author Tel.: +964 -7829090612 Email: mohammed.g.mukhelif@tu.edu.iq

ABSTRACT

Novel Imine were prepared via the reaction of (p-chloro and p-methyl)aniline with (p-nitro and p-methoxy) benzaldehyde and later engaged in a condensation reaction with 2-O-Acetylmalic anhydride in a dioxane as a solvent. The resulting products were found to be 1,3-oxazepine-1,5-diones and their derivatives. FT-IR, ¹HNMR and C.H.N. spectra were used to confirm the structures of products. The antibacterial properties of these compounds were also examined

Keywords:

Seven-Member, Antibacterial Activities, 1,3-Oxazepine, Imine

Introduction:

Oxazepine has a seven-member unsaturated non-homologous ring containing two heteroatoms, oxygen and nitrogen, and five carbon atoms. [1] One sort of pericyclic reaction used to make 1,3-oxazepine was the cycloaddition process. [2] Several studies have found that imported Oxazepine compounds exhibit a wide range of biological actions, including antibacterial [3], antifungal [4], antitumor [5], anti-influenza [6], antianxiety[7], antipsychotic[8], anticonvulsant[9], anticancer [10], They also have anticorrosion properties. [11]

The target of this research is to create novel (1,3) oxazepine 1,5-diones molecules, characterize them, The antibacterial properties of these compounds were also examined.

Material And Methods

The use of all chemicals was done without additional purification and they were all bought from Sigma-Aldrich. The infrared spectra were captured using a Tensor 27 Bruker, Germany spectrometer (spectral range: 4000-600 cm⁻¹). The ¹H NMR spectra were captured using DMSO-d₆ as the solvent on a Bruker Ultershield 400 MHz NMR spectrometer in Germany. The C.H.N analyses were performed by Euro Ea Elemental Analyser.

Procedure for prepare of Schiff's Bases A₁-A₄

A solution of p-methylaniline or p-chloro aniline (0.01 mol) in Ethanol absolute (15 mL) was added to 50 mL round bottom flask containing appropriate aldehyde (0.01 mol). Refluxing the mixture for three hours, After

filtering, the precipitate was re-crystallized from ethanol. Table 1 provides a list of some of

the physical characteristics of compounds prepared




Table (1): Some of the Physical properties of compounds A1-A4

Com. Symb.	Formula	X	Y	Yields		Colour
				Mol.Wt.	%	
A ₁	C ₁₄ H ₁₂ ClNO	Cl	OCH ₃	245.71	74	white
A ₂	C ₁₃ H ₉ ClN ₂ O ₂	Cl	NO ₂	260.68	83	Yellow
A ₃	C ₁₅ H ₁₅ NO	CH ₃	OCH ₃	225.29	71	white
A ₄	C ₁₄ H ₁₂ N ₂ O ₂	CH ₃	NO ₂	240.26	79	Yellow

Procedure for prepare of 1,5-Disubstituted oxazepine

In a condenser-equipped, 100-ml round-bottom flask that has been thoroughly dried, a mixture of equivalent amount (0.001mole) of Schiff's bases A(1-4) and (0.001mole) of 2-O-Acetylmalic Anhydride in anhydrous Benzene (30 ml) was refluxed for (4 hr.). In an ice bath,

the reaction mixture was allowed to cool. separated out some compounds as solid product during cooling, and some other as a Gummy. The product was washed in distilled water. and later dried and recrystallized from benzene. The chemical formula, Molar mass, yield %, melting points and colours, are given in table (2).




Table (2): The structural formula, yield, colors of compounds


Symb.	Formula	X	Y	Yields		M.P. °C	Colour
				Molar mass	%		
B ₁	C ₂₀ H ₁₈ ClNO ₆	Cl	OCH ₃	403.82	76	Gummy	Brown
B ₂	C ₁₉ H ₁₅ ClN ₂ O ₇	Cl	NO ₂	418.79	59	Gummy	Brown
B ₃	C ₂₁ H ₂₁ NO ₆	CH ₃	OCH ₃	383.40	78	Gummy	Brown
B ₄	C ₂₀ H ₁₈ N ₂ O ₇	CH ₃	NO ₂	398.37	68	Gummy	Brown

Results and Discussion

The reaction of anhydride derivatives as an electrophilic reagent with imines (Schiff's

bases) as mild nucleophilic reagents is described in this paper.

Schiff's bases were prepared using an acid catalyzed thermal condensation reaction



The mechanism of imine synthesis has been fully explained in the literature.[12-13]

The structures of imine were confirmed by their FT-IR spectra which showed the disappearance and appearance of the characteristic absorption frequencies (bands) of the principal functional groups. According to the FT-IR spectra, the distinctive absorption frequencies of both (C=O) at (1720-1740) cm^{-1} and (-NH₂) at (3300-3500) cm^{-1} vanished, of

with p-methylaniline or p-chloroaniline and an appropriate aldehyde, according to a well-known procedure.


the primary amine and the aldehyde, respectively. and the occurrence of the stretching absorption bands of the azomethine group (C=N) at (1619-1625) cm^{-1} , in addition to the appearance of stretching absorption of the other groups in the structure of each individual compounds table (3).

Table (3): FT-IR data of imine compounds [A₁-A₄]

Comp.	ν C-H ar. st.	ν C=N st.	ν C=C st.	ν C-N st.	δ (C-H) Bending	Other cm^{-1}
A ₁	3018	1619	1567	1477 1090	726, 838	2962 C-H al. st. 1371 C-NO ₂
A ₂	3099	1624	1597	1487 1008	713, 823	sy. 1340 C-NO ₂ as.
A ₃	3077	1621	1595	1504 1022	724, 835	2971 C-H al. st. 1376 C-NO ₂
A ₄	3078	1625	1596	1514 1005	711, 836	sy. 1339 C-NO ₂ as.

By mixing equivalent amounts of anhydride derivatives in anhydrous Benzene and refluxing under dry conditions, the reaction of

anhydride derivatives with a variety of Schiff's bases was studied. The reaction's plausible reaction mechanism was described as follows:



In this work, the (2+5) polar cycloaddition reaction of Schiff's Bases as mild nucleophile to 2-O-Acetylmalic Anhydride as electrophile in anhydrous Benzene is reported. The structures of the synthesized oxazepine were confirmed by the FT-IR spectra which showed the disappearance and appearance of the characteristic absorption frequencies (bands) of the principal functional groups. The FT-IR spectra showed the disappearance of the characteristic absorption frequencies of

azomethine group (C=N) at (1619-1625) cm^{-1} , and the appearance of the stretching absorption bands of (C=O lactone) at (1750-1765) cm^{-1} , (C=O lactame) at (1673-1711) cm^{-1} in addition to the appearance of stretching absorption of the other groups in the structure of each individual compounds Figure 1 and Figure 2 : IR spectrum of compound B₁, ¹H NMR spectra confirm the structures of synthesized compounds. Figure 3 and Figure 4: ¹H NMR spectrum of B₁₋₄. [14-16]

Table (4): FT-IR data of 1,3-oxazepines derivatives

Comp.	ν C-H ar. st.	ν C=O lactone	ν C=O lactame	ν C=C ar. st.	ν C-N st.	δ (C-H) Bending	Other cm^{-1}
B ₁	2998	1750	1673	1572 1522	1090	739, 858	2962 C-H al.st.
B ₂	3104	1761	1680	1596 1517	1010	739, 851	1400 C-NO ₂ sy. 1341 C-NO ₂ as.
B ₃	3075	1765	1709	1591 1515	1018	718, 837	2992 C-H al. st.
B ₄	3021	1759	1711	1600 1518	1008	739, 842	1400 C-NO ₂ sy. 1344 C-NO ₂ as.

Table (5): Chemical Shift δ (ppm) and (C.H.N.) analysis of 1,3-oxazepines derivatives

Comp	Formula	M.Wt (g/mole)	Chemical Shift δ ppm	C.H.N % Calculated (Found)		
				C%	H%	N%
B ₁	C ₂₀ H ₁₈ ClNO ₆	403.82	(1Hs) δ =8.07 (N-C-H), (8Hm) δ =7.09-7.64 (-C-H _{arom.}), (1Hm) δ = 5.59-5.60 (O-C-H) (3H s) δ = 4.07 (OCH ₃) (2H d) δ = 2.94-2.97 (CH ₂ ring7) (3H s) δ = 1.72 (O=C-CH ₃)	59.49 (58.87)	4.49 (4.27)	3.47 (2.84)
			(1Hs) δ =8.15 (N-C-H), (8Hm) δ =6.88-7.58 (-C-H _{arom.}), (1Hm) δ = 5.19-5.21 (O-C-H) (2H d) δ = 3.66-3.68 (CH ₂ ring7) (3H s) δ = 2.11 (O=C-CH ₃)	54.49 (53.76)	3.61 (2.93)	6.69 (5.59)
B ₃			(1Hs) δ =8.18 (N-C-H),			

	$C_{21}H_{21}NO_6$	383.4 0	(8Hm) δ =6.92-7.18 (-C-H _{arom.}), (1Hm) δ = 6.11-6.14 (O-C-H) (3H s) δ = 4.03 (OCH ₃) (2H d) δ = 3.27-3.29 (CH ₂ ring7) (6Hs) δ =2.68(O=C-CH ₃),(Ar-CH ₃) (1Hs) δ =7.98 (N-C-H), (8Hm) δ =7.09-7.77 (-C-H _{arom.}), (1Hm) δ = 5.56-5.60 (O-C-H) (2H d) δ = 2.92-2.95 (CH ₂ ring7) (6Hs) δ =2.68(O=C-CH ₃),(Ar-CH ₃)	65.79 (63.79)	5.52 (4.93)	3.65 (3.09)
B ₄	$C_{20}H_{18}N_2O_7$	398.3 7	(8Hm) δ =7.09-7.77 (-C-H _{arom.}), (1Hm) δ = 5.56-5.60 (O-C-H) (2H d) δ = 2.92-2.95 (CH ₂ ring7) (6Hs) δ =2.68(O=C-CH ₃),(Ar-CH ₃)	60.30 (60.79)	4.55 (3.93)	7.03 (6.89)




Figure 1: IR spectrum of compound B₁.




Figure 2: ^1H NMR spectrum of compound B₁.

Biological Activity against Bacteria

The heterocyclic derivatives of 1,3-oxazepines have been shown to have antibacterial action against a variety of bacteria, including *E. coli*, *S. aureus*, and *P. aregenosa*. utilizing nutritional agar medium and the well diffusion method. Each substance was suspended in aqueous

solutions at various concentrations (10–100 mg/mL), Solvent blanks were used in all assays against each test organism, and the results are represented as MICs (minimal inhibitory concentrations). The experimental biological data is shown in Table 6.

Table 6: Antibacterial activity data of the heterocyclic derivatives of 1,3-oxazepines

Comp.	<i>E. coli</i>	<i>S. aureus</i>	<i>P. aregenosa</i>
B ₁	22	21	22
B ₂	23	21	19
B ₃	17	19	20
B ₄	20	20	18
Antibiotics	-----		-----
Ampicillin	23	20	21
Vibromycin	24	22	20

Conclusions

In this work, the new substituted 1,3-oxazepine-4,7-dione derivatives were prepared successfully by using various substituted Schiff's bases. The 1,3-oxazepines has been evidenced by spectral analysis. The substances' antibacterial efficacy against several types of bacteria was assessed, and they shown similar activity to that of conventional medications.

References:

1. A- Al-awwadi, N. A. J.; Alsaffee, B. A. H.; Abdulridha, M. M. (2016) "Synthesis and Characterization of Cu(II) and Fe(II) metal complexes of Oxazepine derivative via Schiff base [Fe(HPOHBOT)Cl₂] and [Cu(HPOHBOT)Cl₂]", African Journal of

- Pharmacy and Pharmacology 10(35): 728-736.
2. Al-Hiti, W. F.; Saeed, M. A. (2017) "Synthesis and Characterization of some 3-(2-(6-oxo-1,3-thiazinan-3-yl)-R-)-1,3-oxazepine-4,7-dione and N-Bromoamines 1,3-oxazepine-1,4-dione derivatives", *BEST: International Journal of Humanities, Arts, Medicine and Science* 5(1): 65-74.
 3. Jassim, I.K.; Ali, A. M. (2018) "Synthesis and Studying The Liquid Crystalline Properties for New Heterocyclic Compounds", *Journal of Education and Scientific Studies-Chem Sci.* 12(3):172-183.
 4. Muslim, R. F.; Tawfeeq, H. M.; Owaid, M. N.; Abid, O. H. "Synthesis, Characterization and Evaluation of antifungal activity of seven-membered heterocycles", *Acta. Pharm. Sci.* 56(2):39-57. (2018)
 5. uni, D.; Ranjitha, C.; Rama, M.; Pai, K. S. R. "Oxazepine derivative as an antitumor agent and snail1 inhibitor against human", *International Journal of Innovative Research in Science, Engineering and Technology* 3(8): 15357-15363. (2014)
 6. El-Nezhawy, A. O. H.; Eweas, A. F.; Maghrabi, I. A.; Edalo, A. S.; Abdelwahab, S. F. "Design, Synthesis and Molecular docking of novel Pyrrolooxazepinediol derivatives with anti-influenza neuraminidase activity" *Arch. Pharm. Chem. Life Sci.* 348(11):786-795. (2015)
 7. Mahapatra, D. K.; Shirhare, R. S.; Gupta, S. D. "An oxyolytic activity of some 2,3-dihydrobenzo[b][1,4]-oxazepine derivatives synthesized from Murrayanine-Chalcone", *Asia J. Res. Pharm. Sci.* 8(1): 25-29. (2018)
 8. Naporra, F.; Gobleder, S.; Wittmann, H.; Spindler, J.; Bodensteiner, M.; Berhardt, G.; Hubner, H.; Gmeiner, P.; Elz, S.; Strasser, A. "Dibenzo[b,f][1,4]oxazepiens and dibenzo[b,e]oxepines: influence of the chlorine substitution pattern on pharmacology at the H₁R, H₄R, 5-HT_{2A}R and other selected GPCRs", *Pharmaceutical Research* 113(A):610-625. (2016)
 9. Sahu, M.; Nerka, A. G.; Chikhale, H. U.; Sawant, S. D. "In silico screening, synthesis and pharmacological screening of quinazolinones containing oazepinone ring as NMDA receptor antagonists for anticonvulsant activity: part-I", *Journal of Young Pharmacists* 7(1):21-27. (2015)
 10. Al-Rawi, M.S.; Hussei, D.F.; Al-Taie, A. F.; Al-Halbosiy, M.M.; Hameed, B. A. "Cytotoxic effect of new synthesis heterocyclic derivatives of amoxicillin on some cancer cell lines", *IOP Conf. Series: Journal of Physics: Conf. Series* 1003(012012):1-7. (2018)
 11. Hamak, K.F.; Eissa, H.H.(2013) "Synthesis characterization, biological evaluation and anti-corrosion activity of some heterocyclic compounds Oxazepine derivative from Schiff bases", *International Journal of ChemTech Research* 5(6):2924-2940.
 12. William, H.; Brown, B.L.; Iverson, E.V.; Anslyn, C.S.F. "Organic Chemistry", 7th Ed., WADSWORTH CENGAGE Learning. (2014)
 13. McMurry, J. (2016) "Organic Chemistry", 9th Ed., CENGAGE Learning.
 14. L.D.S.Yadav, *Organic Spectroscopy*, Springer Science,Business Media Dordrech, Springer-Science.Business Media, B.V. (2005).
 15. Donald L. Pavia, Gary M. Lampman, George S.& James R. Vyryan, *Introduction to spectroscopy*, 4th Ed., Brooks\ cole cengage Learning, pp 16-104, (2009).
 16. Robert M. Silverstein, Francis X. Webster, David J. Kiemle, *spectrometric Identification of Organic compounds*, 7th Ed, John- Wiley & Sons, INC., pp. 72-126, (2005).