



# Analysis Of Combinatorial Aspects Of Random Events

**Shingis Maykhanovich Urmanov**

Chirchik State Pedagogical University  
[urmanovshingis@gmail.com](mailto:urmanovshingis@gmail.com)

## ABSTRACT

This article examines the basic concepts of combinatorics in the context of probability theory. The types of random events and methods for their description using combinatorial approaches are analyzed. Particular attention is paid to the classification of events and methods for calculating their probabilities.

**Keywords:**

Test, event, probability theory, probability

**The Introduction.** A significant part of mathematics, probability theory is devoted to the investigation of random occurrences and occurrences that are not predetermined. On the other hand, combinatorics offers instruments that may be utilized for the purpose of examining the numerous permutations and combinations of components that are contained inside these random events. The categorization of random occurrences serves as an example for the purpose of this article's investigation of the connection between combinatorics and probability theory.

Conflicts and occurrences take place. A trial is an observation of a random phenomena that can lead to the occurrence of a variety of outcomes, according to the theory of probability. In the context of a trial, an event refers to a particular outcome or group of events.

A procedure or experiment that is related with random events and from which the outcomes cannot be anticipated in advance is referred to as a trial. Instances of trials include things like flipping a coin, rolling dice, drawing a card from a deck, and other similar activities.

It is possible that each experiment will result in one of several different results. When a trial is carried out, there is the potential for a number of different outcomes, which may be referred to as events. When it comes to events, they can either be simple, meaning they consist of a single consequence, or composite, meaning they consist of several outcomes. A few examples of events are obtaining heads on a coin toss, receiving an even number on a die roll, and other similar types of occurrences.

For instance: Tossing a coin is the first trial. - The results are Heads and Tails. Heads and Tails are the events. Throwing a die with six sides is the second trial.

- The results are as follows: 1, 2, 3, 4, 5, 6. Even numbers (2, 4, 6), numbers that are more than 4 (5, 6), and particular numbers, such as 3, are examples of events. The third step is to select one card from a typical deck of 52 playing cards.

- The results include a total of 52 distinct cards, such as the ace of hearts, the king of spades, the queen of diamonds, and so on. Drawing a black card, drawing a spade card, drawing an ace, and other events are examples

of events. Principles of significance: At the elementary level, an event is defined as one that cannot be subdivided into simpler events. Consider the case of rolling a three on a dice. An event that is comprised of many elementary events is referred to as a composite event. Take, for instance, the act of rolling a die with an even number (2, 4, 6).

Certain occurrence refers to an event that consistently takes place whenever a trial is carried out. For instance, using a die with six sides and rolling a number between one and six.

An instance that never takes place during the course of a trial is referred to as an impossible occurrence. Consider the case of rolling a seven on a die with six sides. The probability of an event is defined as the ratio of the number of positive outcomes to the entire number of potential outcomes of the trial, assuming that all options are equally likely. In other words, the probability of an event A depends on the number of favorable outcomes:

$$P(A) = \frac{\text{number of favorable outcomes}}{\text{total number of outcomes}}$$

where  $P(A)$  — probability of event A.

These ideas and procedures form the foundation of probability theory and are utilized extensively in a variety of professions, including but not limited to statistics, economics, engineering, science, and others like these.

On the basis of their characteristics and the way in which they are connected to the results of the test, random occurrences may be classified into three primary categories:

There are several kinds of random occurrences. Random occurrences can be categorized according to their structure and the way in which they are connected to the results of the test. Among the most common kinds of random occurrences are:

#### Simple Occurrences (a)

A single outcome of a trial is referred to as a simple event, sometimes known as an elementary event. It is not possible to break them down into more manageable occurrences.

For instance:

When you flip a coin, the most straightforward outcomes are either getting heads or getting tails.

A simple event is the occurrence of the numbers 1, 2, 3, 4, 5, or 6 when a die with six sides is thrown.

#### c) Events Together Made Up

Events that are composite, also known as complicated events, are those that involve more than one outcome of a trial. The union of these events is represented by them, which encompass a number of elementary occurrences collectively.

For instance:

In the process of tossing a die with six sides, the occurrence known as "getting an even number" is comprised of the results 2, 4, and 6.

The outcomes of the occurrence known as "drawing a black card" from a typical deck of cards include all of the spades and clubs in the deck.

b) Other Related Occasions What are known as additional events, sometimes known as opposing events, are events that encompass all outcomes that are not included in the event that is being discussed. In the context of an event A, an extra event is represented by the symbol  $- A$ , which encompasses all outcomes that are not included in the overall event A.

An example of this would be the following: while tossing a coin, getting tails is an additional event to obtaining heads. while throwing a six-sided die, receiving an odd number (1, 3, 5) is an additional event to getting an even number (2, 4, 6).

#### (d) Unimaginable Occurrences

During a particular trial, "impossible events" are events that are not feasible to occur. It is always the case that the probability of an impossible event is zero.

As an illustration, it is not feasible to get the result of "getting seven" on a die with six sides, as the die only has numbers ranging from one to six.

Due to the fact that there are only four suits in a conventional deck of cards, the occurrence of "drawing a card of the eighth suit" is not capable of occurring.

#### e) Specific Occurrences

Within the context of a trial, there are some occurrences that are guaranteed to take place. Without fail, the likelihood of a particular occurrence is always one.

One example is that when a six-sided die is rolled, the event of "rolling a number from 1 to 6" is guaranteed since the die only has these six sides instead of any other number.

The occurrence of "drawing a card from a deck of 52 cards" is guaranteed at all times, as it is always the case that one card is pulled from this deck.

#### f) Events that are Exclusive and Private

In the context of a single trial, joint events are defined as occurrences that potentially take place concurrently.

As an illustration, when two dice are rolled, the event "rolling an even number on the first die" and the event "rolling a three on the second die" are considered to be joint events since they both have the potential to take place at the same time.

In the context of a single trial, exclusive events are defined as occurrences that cannot come about simultaneously.

As an illustration, when a single coin is rolled, the outcomes of "rolling heads" and "rolling tails" are mutually exclusive. This is because only one of these outcomes may take place during a single examination.

In a variety of experiments and circumstances, these categories make it possible to conduct a more in-depth investigation of random occurrences and the probability associated with them.

Within the realm of probability theory, random occurrences are an essential component in the modeling and analysis of random incidents. It is feasible to categorize these occurrences according to a variety of criteria, and they reflect subsets of all the conceivable outcomes the study may produce. The primary categories of random occurrences and the characteristics that distinguish them will be discussed in this article.

Complex occurrences occurring. It is possible to express a complex event as a union or intersection of numerous elementary events. A complex event is made up of several basic events.

#### Examples:

- When a dice is thrown, the event "an even number came up" (2, 4, or 6) is an example of a

complicated event that is composed of simple events.

In the process of tossing two coins, the event known as "both coins come up heads" is the result of the intersection of two fundamental events: the first coin lands on its head, and the second coin lands on its head.

The events are compatible. If it is possible for two events to take place at the same time, then we say that they are compatible. This means that their intersection is not empty.

#### Examples:

- When two coins are flipped, the occurrences of "getting heads on the first coin" and "getting tails on the second coin" are compatible with one another since they can occur concurrently.

The "getting a pair" and "getting three of a kind" occurrences in the card game "poker" are also compatible with one another. This is due to the fact that it is possible to obtain a combination that consists of both a pair and three of a kind.

Events that are incompatible. Two events are said to be incompatible if they are unable to occur concurrently, which means that their intersection is not filled by any events.

Example: - When tossing a die with six sides, the events "3" and "4" are incompatible with one another since the same trial cannot give two different numbers at the same time.

-! The events "product is in good working order" and "product is defective" are incompatible with one another in the context of product testing. This is due to the fact that a product cannot be in both excellent working order and faulty positions at the same time.

Carry out the events. When all of the conceivable outcomes of a trial are covered by the union of a set of events, that set of events is said to be complete. In other words, it contains all of the possible elements that make up the space of elementary events.

To illustrate, when a coin is flipped, the set of events known as "heads" and "tails" is considered to be complete. This is because the set encompasses all possible outcomes that can occur from flipping a coin.

-! It is also true that the set of occurrences "a number from 1 to 6" is comprehensive when

it comes to the act of tossing a die because it encompasses all of the potential outcomes of a trial.

Not affiliated with any other event. To say that two occurrences are independent is to say that the likelihood of their happening together is equal to the product of the probabilities of each of the events individually.

In a formal sense, the independence of events A and B is shown by the equation  $P(A \cup B) = P(A) * P(B)$ .

Some examples include the following: - When two coins are tossed, the result of the first toss does not influence the result of the second toss. The outcomes of "heads on the first coin" and "heads on the second coin" are therefore considered to be separate events.

The occurrences of "winning on the first ticket" and "winning on the second ticket" are considered to be independent in the context of a lottery, in which the draw of one ticket does not influence the likelihood of winning the other ticket.

The events are complementary. Events that are complementary to one another are those that, when taken collectively, constitute the whole set of all outcomes that are possible. It is necessary for one of them to take place in the event that the other does not take place.

In a formal sense, if A is an event, then its complementary event  $A^c$  is the opposite event, where the probability of A plus the probability of  $A^c$  equals 1.

For instance:

It is important to note that the event "heads" and the event "tails" are complimentary when flipping a coin.

There is a complimentary relationship between the occurrences "test positive" and "test negative" when it comes to testing for an illness.

Finally, the Conclusion. Combinatorics is an important part of the analysis of random occurrences because it enables us to define and compute the probability of a number of different possibilities for the outcomes of trials. By gaining an understanding of the fundamental combinatorial procedures, we are able to investigate probability theory and its

applications in a variety of domains in a more in-depth and accurate manner.

## References:

1. Уильям Феллер (William Feller) "Введение в теорию вероятностей и ее приложения". Издательство "Наука" 1984 (перевод на русский язык)
2. Castro, M.J., Garcia-Rodriguez, J.A., Gonzalez-Vida, J.M., Macias, J., Pares, C., VazquezCendon, M.E., 2004. Numerical simulation of two-layer shallow water flows through channels with irregular geometry. *J. Comput. Phys.* 195, 202-235.
3. Chang, T.-J., Chang, K.-H., Kao, H.-M., 2014. A new approach to model weakly nonhydrostatic shallow water flows in open channels with smoothed particle hydrodynamics. *J. Hydrol.* 519, 1010-1019.
4. А. Н. Ширяев "Теория вероятностей и математическая статистика". Издательство: "Физматлит" 2004
5. Л. В. Богоявленский "Комбинаторика". Издательство: "Наука" 1986
6. Кеннет Х. Розен (Kenneth H. Rosen) "Дискретная математика". Издательство: "Мир" 2001.
7. Raupova, M. H., & Xasanova, M. F. (2024). PERFORMING COMPLEX MATHEMATICAL CALCULATIONS IN PYTHON USING THE SYMPY LIBRARY. *European Journal of Interdisciplinary Research and Development*, 25, 73-78.
8. Raupova, M. H., Xasanova, M. F., & Qurbanqulova, S. B. (2024). PYTHON DASTURLASH TILINING SYMPY VA SCIPY KUTUBXONALARIDAN FOYDALANGAN HOLDA INTEGRALGA DOIR MASALALARINI HAL QILISH. *PEDAGOG*, 7(2), 523-528.
9. Раупова, М. Х., & Урманов, Ш. М. (2024). ИЗМЕНЕНИЯ ПЕДАГОГИЧЕСКОГО ПРОЦЕССА В СОВРЕМЕННОЙ ШКОЛЕ: ВЫЗОВЫ И ПЕРСПЕКТИВЫ. *PEDAGOG*, 7(2), 515-522.

10. Raupova, M. H. (2024). USING QUANTUM ALGORITHMS IN CRYPTOGRAPHY. *Web of Technology: Multidimensional Research Journal*, 2(3), 9-14.
11. Mukhamedieva, D. T., & Raupova, M. H. (2024). Model of biodiversity and plant sustainability based on quantum variational optimization. In *E3S Web of Conferences* (Vol. 498, p. 02007). EDP Sciences.
12. Mukhamedieva, D. T., & Raupova, M. H. (2024). Model for forest ecosystems based on quantum optimization. In *E3S Web of Conferences* (Vol. 498, p. 02006). EDP Sciences.
13. Panjiyeva, G., & Raupova, M. (2024). A DEEPER UNDERSTANDING OF THE CONCEPT OF A FUNCTION IN A PYTHON PROGRAM. *PEDAGOG*, 7(4), 514-518.
14. Jamolov, B., & Raupova, M. (2022). SOME REMARKS ON EQUATIONS OF THE HYPERBOLIC TYPE WITH PARTICULAR DERIVATIVES. *Science and Innovation*, 1(6), 301-316.