



Agrochemical Properties Of Pomegranate Crops

Turdimetov Shaxobiddin Muxitdinovich

Doctor of Biological Sciences(DSc), professor of the Department of Soil Science and Reclamation at Gulistan State University.
e-mail turdimetov1970@mail.ru

G'oibova Sevara Ayup qizi

Gulistan state university, student.
e-mail sevaragoibova911@gmail.com

ABSTRACT

Presents reviews of the work of local and foreign scientists on the importance of studying the agrochemical properties of the soil in determining the rate of pomegranate fertilization, and the pomegranate's demand for fertilizer. Information about the soil-climatic conditions of the research object is given and the results obtained on agrochemical properties are described. Recommendations on fertilization periods and norms are presented, taking into account the agrochemical properties of the soil.

Keywords:

Pomegranate, Soil, Nitrogen, Phosphorus, Potassium, Supply Of Nutrients, Stratified Fertilizing.

Enter. It is important to know the level of supply of nutrients in the soil when determining the rates of fertilization of agricultural crops. Because in addition to the biological demand of crops for nutrients, the amount of nutrients in the soil is also taken into account in fertilizing. According to the amount of nutrients in the soil, it is possible to use fertilizers in a stratified manner.

In Turkey, M.Guneri, M.Yildiztekin, A.L.Tuna, I.Yokas [1] were emphasized the importance of fertilizing pomegranate with potassium and calcium, and considered that it is necessary to take into account its amount in the soil. In research, the effect of potassium and calcium elements on the quality and amino acid content of the fruit of the pomegranate plant was studied. Pomegranate fed with enough potassium and calcium increased fruit weight, juice yield and skin thickness, and improved fruit quality.

In India, scientists of the Department of Fruit Production of the Punjab Agricultural

University, P.P.S.Gill, M.Kumar, N.P.Singh, W.S.Dhillop [2] conducted an experiment using NPK fertilizers in pomegranate crops at different rates. Application of 300:50:100 g of NPK per plant was found to be the most optimal option, and plant growth and productivity were the highest in this option. The color of the fruit, the number of grains and the thickness of the shell were better in the plant well fed with potassium compared to the control variety. It is emphasized that the agrochemical parameters of the soil are important in determining the application dose of fertilizers.

S.Davarpanah et al. [3] found that fruit cracking was reduced by 26-52% compared to control when pomegranate crops were moderately fertilized with calcium in Iran. In addition, fruit yield and quality were good when fertilized in moderation. This is due to the increase in the activity of amino acids and the improvement of other biochemical processes.

V.M.Prasad [4] studied the effect of nitrogen, potassium and boron on growth, flowering, fruiting and fruit quality of pomegranate plant in 2019-2020. Experiments were conducted in 10 variants and 3 repetitions. During biometric observations, pomegranate growth and fruiting processes were fully controlled. The agrophysical, agrochemical and biological properties of the soil are improved under optimally fertilized conditions.

A lot of experiments have been carried out on determining the norms of pomegranate fertilization, taking into account the agrochemical properties of the soil, the effect of fertilizers on productivity [5-7]. According to the results of experiments, the rates and periods of fertilizing are different in different regions and differ from each other according to the soil and climatic conditions. The reviews on pomegranate fertilization given above also contain information on the use of calcium fertilizers for fertilization. The amount of calcium in our soils is sufficient, even excess. Therefore, it is important to determine the fertilization standards taking into account the soil and climate conditions.

Research methods

Special experiments were carried out to calculate the fertilization rates taking into account the agrochemical properties of the soil. Soil samples were taken and analyzed. The amount of humus, total nitrogen, phosphorus and potassium, nitrate nitrogen, mobile phosphorus, exchangeable potassium, carbonates, and gypsum in the soil was determined. The absorption capacity of the soil and the content of cations were also determined. On the basis of these data, recommendations on the rate of pomegranate fertilization were developed.

Results and discussions

The experimental area planted with pomegranate is located in the central part of the Mirzachol oasis and consists of irrigated gray-meadow soils. The amount of humus in the soils of the studied area in most cases does not reach 1%. This indicates that these soils are low in humus. The reason for the low content of humus and total nitrogen is explained by the reduction of organic residues as a result of soil salinization, and the low ability of nitrification.

Agrochemical indicators of the soil are one of the properties that determine its productivity. The amount of humus in the soil is higher in the surface layer (Table 1).

Table 1.
Agrochemical indicators of grassland soils.

Section №	Depth, sm	hummus %	SO ₄ gypsum, %	SO ₂ carbonates, %
24-1	0-30	1,139	0,173	9,17
	30-50	0,696	0,226	8,57
	50-100	0,528	0,207	9,14
24-2	0-30	0,945	0,182	9,15
	30-50	0,622	0,174	9,18
	50-100	0,345	0,192	10,16

In the soil hummus quantity every cut into pieces too don't drive in a row high to the indicator owner and 0.945-1.139 percent organization will do. Soil under the knife is 0.622-0.696%, down side while his amount up to 0.345 percent decreased.

Carbonates in the research area are distributed almost evenly on the soil profile and its amount is 8.5-10.0%. This indicates the weak alkalinity of these soils. The amount of gypsum ranges from 0.173% to 0.207%. An increase in its amount causes deterioration of soil properties.

The amount of total nitrogen depends on the amount of humus, and its amount is also high in the arable layer (Table 2).

24-1- equal to 0.072 percent in the 0-50 sm layer of the section. In both sections, its

amount has decreased. The amount of total phosphorus reaches 0.3 percent in the 24-2 section. Also, the total amount of potassium was higher in the 24-2 section and increased to 1.44%.

Table 2.
Amount of total and mobile nutrients in grassland soils.

Section №	Depth, sm	General, %			Mobile mg/kg		
		Nitrogen	Phosphorus	Potassium	N-NO3	P2O5	K2O
24-1	0-30	0.072	0.185	0.57	17.4	27.0	113.2
	30-50	0.046	0.145	0.47	13.5	13.0	96.3
	50-100	0,039	0,128	0,42	11,0	10,4	89,1
24-2	0-30	0,056	0,288	1,12	20,2	15,3	280,2
	30-50	0,038	0,254	1,14	15,4	10,9	170,1
	50-100	0,032	0,212	1,44	12,6	10,4	120,6

The amount of nitrate nitrogen was 20.2 mg/kg in the arable layer of section 24-2, its amount decreased to 11.0 mg/kg in the 50-100 sm layer of section 24-1. The amount of mobile phosphorus is 27.0 mg/kg in the arable layer of section 24-1, 15.30 mg/kg in section 24-2, and 10.4 mg/kg downward. This indicates that these soils are poor in mobile phosphorus. The main reason for this is that gray soils are rich in calcium carbonates and have a high probability of forming calcium phosphates. The amount of mobile potassium is distributed in proportion to the amount of total potassium. Its amount reaches 280.2 mg/kg. Since the parent rocks of gray soils consist of loess, the soils are well supplied with mobile potassium compared to other soils.

In order to determine the rate of fertilization of the pomegranate crop, it is necessary to have complete information about the soil properties. In saline soils, the rate of fertilizers is slightly increased. There is enough information about the properties of the soils of the research object, their improvement [8-17]. The task now is to organize the fertilization of the pomegranate crop on a scientific basis based on this information.

Conclusion

The analysis of the agrochemical properties of the soil of the research object shows that the soil is low in nutrients and these

parameters must be taken into account when determining the fertilization rates. In addition, it is necessary to use fertilizers differentially, taking into account the age, variety, and planned yield indicators of the pomegranate crop.

References:

1. Guneri M., Yildiztekin M., Tuna A.L., Yokas I. 2016. Calcium and potassium fertilization on antioxidant capacity and the fruit quality of 'hicz' pomegranates (*Punica granatum*L.). *Bulg. J. Agric. Sci.*, 22: pp. 591–599.
2. Gill P.P.S., Kumar M., Singh N.P., Dhillon W.S. Studies on macronutrient fertilization in pomegranate under sub-tropical plains. *J.Hortl. Sci.*Vol. 8(2):172-175, 2013.
3. Davarpanah S. et al. Foliar calcium fertilization reduces fruit cracking in pomegranate (*Punicagranatum*cv. Ardestani). *Scientia Horticulturae* 230 (2018) 86–91.
4. Kuldeep and Prasad V.M. 2020. Effect of Nitrogen, Potash and Boron on Plant Growth, Flowering, Fruit Drop and Fruit Setting of Pomegranate (*Punica granatum*) cv. Bhagwa. *Int.J.Curr.Microbiol.App.Sci.* 9(11): 2181-2185.
5. Dhillon W.S., Gill PPS, SinghN.P. Effect of Nitrogen, Phosphorus and Potassium

- Fertilization on Growth, Yield and Quality of Pomegranate 'Kandhari'. Article in Horticultural Actae. March 2011 DOI: 10.17660/ActaHortic.2011.890.45.
6. Chidananda, G., BS Shivakumar, Sarvajna B. Salimath, R. Girish and Chaitanya, H.S. Bhagwa. Int.J.Curr.Microbiol.App.Sci. 9(10): 2885-2
 7. Priya, B Kurubar, Ashok AR, Ramesh H, Udaykumar G, Umesh MR and Rajkumar RH(2022). Effect of Nitrogen, Phosphorus and Potassium Fertilization on Yield and Quality of Custard Apple (*Annonas squamosal* L.) Cv. Balanagar. Biological Forum – An International Journal, 14(2a): 84-88.
 8. Turdimetov Sh., Esonboyeva N. Land reclamation condition of Mirzaabad district. International scientific and educational electronic journal "EDUCATION AND SCIENCE IN THE XXI CENTURY". Issue No. 37 (vol. 3) (April, 2023). pp. 67-73.
 9. Turdimetov Sh., Esonboyeva N Mirzaobod tumani gidromorf tuproqlarinig xossalari. Eurasian Journal of Technology and Innovation. Volume 1, Issue 5, May 2023. pp 81-85.
 10. Turdimetov Sh.M., Mirsharipova G.K., Botirova L.A., Mustafakulov D.M., Abdujalilova A.X. Impact of legume crops on the agrochemical and agrophysical properties of soil in Mirzachol conditions // – «Journal of Critical Reviews». ISSN – 2394-5125. VOL 7, – ISSUE 17, – 2020. – pp. 2220-2234.
 11. Turdimetov Sh., Khudoyberdiyeva Z., Tadjibayev A. Quality Assessment of Gypsum Soils of Mirzachol Oasis. Journal of Population Therapeutics and Clinical Pharmacology, 2023. 30(12), pp. 295-301.
 12. Turdimetov Sh., Urazalieva M., Esonboeva N. PROPERTIES AND QUALITY ASSESSMENT OF HYDROMORPHIC SOILS OF MIRZACHOL OASIS. Eur. Chem. Bull. 2023,12(Special Issue 1, Part-B), 3992-4000.
 13. Nomozov X.Q., Turdimetov Sh.M. Ozbekiston tuproqlari va ularning evolutsiyasi. Darslik. -T.: «Fan vatexnologiya», 2016, 256 bet.
 14. Turdimetov Sh., Musurmanova M. Properties of Soils located in different Geomorphological Conditions. American Journal of Agriculture and Horticulture Innovations. Volume 02 Issue 11-2022. pp 01-06.
 15. Batirov X.A., Turdimetov Sh., Nurillaeva R.B. Ecological Role of Different Siderate Crops in Improving Soil Properties. American Journal of Agriculture And Horticulture Innovations. Volume 03 Issue 011-2023. pp 01-06.
 16. Shaxobiddin Muxitdinovich Turdimetov, Ilkhomjon Abdulmanzurovich Abdurakhmonov, Laziza Akhmatovna Botirova, Ihtiyor Yakubovich Zikirov, Sevara Mavlonovna Ashiralieva. Soil Quality Assessment Principles for Vegetable Crops. Annals of R.S.C.B., ISSN:1583-6258, Vol. 25, Issue 6, 2021, Pages. 9944 - 9952 Received 25 April 2021; Accepted 08 May 2021.
 17. Turdimetov Sh.M., Sunnatova D. How plant peas affect soil's agrochemical properties // «Modern ecology state environment and scientific practical aspects rational of natural resources problems of agroecology»International Scientific and Practical Conference. – с. Соленое Займище. – 2017. – pp. 779-781.
 18. G'oibova Sevara Ayup qizi "ANOR HAQIDAGI DASTLABKI DA'LILLAR VA ARXEOLOGIK MA'LUMOTLAR" DEVELOPMENT AND INNOVATIONS IN SCIENCE, 1(8), 69-71, 9 june 2022 <https://doi.org/10.5281/zenodo.6629514>.