Eurasian Medical Research Periodical



## Evaluation of Neurotic Disorders in the Post-Covid Period and Treatment Tactics

Tilavov M.T.,	Bukhara State Medical Institute
Kuchkorov U.I	Bukhara State Medical Institute
Barzhakova G.R.	Bukhara State Medical Institute

BSTRACI

Coronavirus infection (COVID-19) is an infectious disease characterized by a relatively high contagiousness and the likelihood of developing severe complications in the form of acute respiratory distress syndrome, acute respiratory and multiple organ failure. The pathogenesis of coronavirus infection has not been studied enough, it has been spreading around the world for the second year. The virus is neurotropic and can directly damage brain structures such as the limbic system and hypothalamus [6].

**Keywords:** 

respiratory distress syndrome

Coronavirus infection Relevance. (COVID-19) infectious disease an characterized by relatively high a contagiousness and the likelihood of developing severe complications in the form of acute respiratory distress syndrome. acute respiratory and multiple organ failure. The pathogenesis of coronavirus infection has not been studied enough, it has been spreading around the world for the second year. The virus is neurotropic and can directly damage brain structures such as the limbic system and Among hypothalamus [6]. the complications after an illness, a significant proportion of patients may develop cognitive disorders: impaired memory, attention, concentration, or speed of thought processes [2,4]. During the study, the authors found that half of patients with coronavirus infection experience a serious level of anxiety and 40% fear that they themselves or their loved ones can become seriously ill with COVID-19 and die [1,5]. Psychopathological symptoms in COVID-19 were as follows: anxiety - 12-20%, depression - 15-25%, insomnia - 8% and

traumatic stress - 35-49% [3]. Based on these data, neurosis-like disorders in the post-COVID period are of great interest and require the development of treatment tactics.

**The purpose**. Systematization of information about neurosis-like disorders in the post-COVID period and improvement of the quality of life of patients.

**Research objectives.** The objective of the study is to study the manifestation of neurosis-like disorders and substantiate new directions of differentiated therapeutic tactics.

Material and research methods. Neurosis- like disorders were studied in the post-COVID period in 46 patients. The average age of the patients was 45 years. All patients were divided into two groups: the first group received basic therapy with nootropic drugs, and the second group, in addition to basic therapy with nootropic drugs, was prescribed the antidepressant rexetine (paroxetine) at a dose of 20 mg 1 time per day for 3 months. After 3 months, the effectiveness of treatment of patients in both groups was evaluated.

Results and discussions. Rexetin was administered at a dose of 20 mg in the morning for 30 days. The condition of patients was assessed before taking the drug (visit 1) and after 15 days (visit 2) and 30 days (visit 3) of treatment. To evaluate the effectiveness of the correctionemotional-volitional sphere used a point scale of subjective characteristics, a

personal and situational scale of anxiety and depression. Each patient received a complex-individual approach. When assessing personal and situational anxiety on the Spielberg-Khanin scale in the post-COVID period, a high level of anxiety prevailed in situational anxiety, and an average level of anxiety prevailed in personal anxiety.

ISSN: 2795-7624

Table 1
Spielberg-Khanin scale of personal and situational anxiety in the post-COVID period before treatment

Type of anyiety	Low Anxiety			level of High level of anxi		of anxiety
Type of anxiety	A1 0/		anxiety		A1 0/	
	Abs.	%	Abs.	%	Abs.	%
situational anxiety	12	26.1	14	30.4	20	43.5
Personal anxiety	10	21.7	24	52.2	12	26.7

After treatment on the 30th day, the indicators in both groups on a scale of situational andpersonal anxiety with a high level of anxiety

turned into a low level of anxiety. Most patients who underwent COVID-19 experienced severe asthenia, anxiety, depression and fear of death.

Table 2 Chanin scale of personal and situational anxiety in the n

Spielberg-Khanin scale of personal and situational anxiety in the post-COVID period after treatment

	Low Anxiety		Average	level of	High level o	of anxiety
Type of anxiety			anxiety			
	Abs.	%	Abs.	%	Abs.	%
situational anxiety	26	56.5	11	23.9	9	19.6
Personal anxiety	29	63.1	9	19.5	8	17.4

Among hospitalized patients, anxiety, confusion, depressed mood and insomnia, as well as impaired attention and memory, were observed in approximately 35%.

Table 3 Scale for assessing the level of reactive and personal anxiety Ch.D. Spielberg and Yu.L. Khanina

Patient group	Test by C.D. Spielberg and Yu.L. Khanin			
	Before treatment	After treatment		
Comparative group	44.6±0.7	31.4±0.7*		
Main group	46.5±0.8	19.7±0.5*,**		

Note: \* - significantly compared to before treatment P<0.001; \*\* - significant compared to control P<0.001

Neurosis-like disorders led to a decrease in the quality of life and a violation of the stability of work. About 60% of patients suffered from insomnia, approximately 20% had suicidal thoughts. In 50% of patients, subdepression was observed, 55% - anxiety, about 60% - psychosomatic symptoms.

Table 4
The severity of asthenic disorders in the post-COVID period

	The severity of asthenic disorders			
Syndrome options	Moderately	Pronounced	Total	
	pronounced			

Total

Astheno-depressive Astheno-hysterical

Astheno-obsessional

Astheno-hypochondriac

4

46

We observed a distinct increase in astheno-adynamic symptoms from 7-12 days of treatment. Against the background of these disorders, a headache appeared, which had a classic "neurasthenic" color. Patients became incapable of emotional, intellectual and physical stress, emotional lability, sensitivity, and a tendency to respond inadequately to all kinds of psycho-traumatic influences intensified. The development of asthenia and emotional stress was characterized by cyclicity. More often in the morning, within 6-12 hours, against the background of severe hypothymia, it arose and quickly covered the entire emotional sphere.

Asthenic with personality changes

According to the results of objective and subjective data, the symptoms of disturbance of the emotional-volitional sphere were noted already on the 3rd day of the disease and persisted even 30 days after treatment. Evaluation of mood according to the results of

the Hamilton test and asthenia (according to the Apathy and Asthenia Scale) was observed in both groups. The main symptoms of asthenia were observed in patients in the form of a change in appetite - 34%, absent-mindedness - 32%, mood instability - 22%, increased excitability with subsequent exhaustion - 29% and increased fatigue - 22.5%.

5

7

2

3

2

1

20

In the combined treatment group (notropes rexetine). and the average improvement in the condition was superior to the basic therapy group. According to the Apathy and Asthenia Scale, in the combined treatment group, compared with the basic therapy group, there was a significantly greater decrease in the severity of such symptoms of asthenia, absent-mindedness, mood instability, increased excitability followed by exhaustion, and increased fatigue.

Table 5
Indicators criteria for quality of life in the post-COVID period

4

7

3

5

4

3

26

Quality of life criteria	Notropes (	(n=24)	Notropes (n=22)	+ rexetine
quanty of the officeria	1st day	30th day	1st day	30th day
General health	46.7±2.3	54.6±1.3	47.4±4.6	69.7±2.3
Physical activity	45.9±3.9	52.1±3.4	46.1±3.8	70.9±3.9
Functioning related to physical condition	48.9±2.7	55.4±2.1	49.6±3.6	68.9±2.7
Functioning related to emotional state	54.3±2.4	59.4±2.8	53.2±4.8	74.3±2.4
Social functioning	55.1±3.2	60.7±2.5	54.2±3.7	68.1±3.2
Pain intensity	56.7±3.1	59.8±1.3	56.8±4.5	66.7±3.1
vital activity	46.6±2.9	52.0±1.4	47.5±3.5	68.6±2.9
Psychological health	49.5±2.8	53.5±1.2	48.3±4.4	65.5±2.8

ISSN: 2795-7624

The evaluation of the therapeutic efficacy of rexetine in the post-COVID period was carried out on the basis of the Hamilton Scale (HDRS). With the use of rexetine against the background of improved mood, a more pronounced regression of emotional disorders was noted. In patients of the main group, pronounced sedation occurred, emotional disturbances were smoothed out, and there was no increase in fear and anxiety in the evening. Along with a mild sedative effect, in the first days of treatment, rexetine caused a positive trend in relation to depression. Patients perked up, made real plans for work.

A significant improvement was noted in comparison with the control group, not only in scales for assessing depression, but also in assessing cognitive functions. Before treatment with rexetine in the main group, the severity of depressive disorder in the main group was 25.6±5.4 points. By the end of the course of therapy, the mean group scores on the Hamilton scale improved significantly, reaching 7.9±1.3 points. In the placebo group, this change was less significant: 24.4±3.8 and 18.9±1.7 points, respectively. Up to 95.6% of clinically stable patients with COVID-19 have psychological problems and symptoms of stress disorders. Psychological distress in the form of depression, hopelessness and nervousness was found in 52% of patients. A statistically significant significant improvement in performance was observed only after 30 days.

**Conclusion.** Treatment of patients who have undergone COVID-19 and have neurosis-like disorders should include correction and therapy of the emotional state to improve the quality of life of patients. Coronavirus infection is a risk

factor for recurrence of neurosis-like disorders. Appointments of the drug Reksetin are a priority method for the correction of emotional-volitional and neurosis-like disorders. The use of antidepressants and nootropics is the most effective in the treatment of somatic and neurosis-like disorders.

## Literature

- 1. Mosolov S.N. "Mental health problems in the context of the COVID-19 pandemic" // Journal of Neurology and Psychiatry. S.S.Korsakov. 2020. V. 120. No. 5. C. 7-15.
- 2. Bao Y., Sun Y., Meng S. et al. 2019-nCoV epidemic: address mental health care to empower society // Lancet. 2020 Vol. 395. P. e37-e38.
- 3. Lai J, Ma S, Wang Y, et al. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. JAMA Netw Open 2020; 3:e203976.
- 4. Rossi R, Socci V, Pacitti F, et al. Mental Health Outcomes Among Frontline and Second-Line Health Care Workers During the Coronavirus Disease 2019 (COVID-19) Pandemic in Italy. JAMA Netw Open 2020; 3:e2010185.
- 5. Schwartz BJ. New Poll: COVID-19 Impacting Mental Well-Being: Americans Feeling Anxious, Especially for Loved Ones. APA News releases. March 25, 2020.
- 6. The COVID-19 Effects of on Hypothalamus: Is it Another Face of SARS-CoV-2 May Potentially That Control the Level of COVID-19 Severity?.2020.